

Cost Drivers of Cancer Care: A Retrospective Analysis of Medicare and Commercially Insured Population Claim Data 2004-2014

Prepared by

Kathryn Fitch, RN, MEd Principal and Healthcare Management Consultant

Pamela M. Pelizzari, MPH Healthcare Management Consultant

Bruce Pyenson, FSA, MAAA Principal and Consulting Actuary

Commissioned by the Community Oncology Alliance

TABLE OF CONTENTS

EXECUTIVE SUMMARY	3
BACKGROUND	5
FINDINGS Cancer population characteristics: 2004-2014 Spending on cancer <i>Components of cancer care</i> <i>Cost characteristics for specific cancer types</i> Chemotherapy site of service <i>Modeling the cost implication of infused chemotherapy site of service shift</i>	7 9 12 15 18 22
DISCUSSION AND CONCLUSIONS	25
APPENDIX A: KEY DATA SOURCES	26
APPENDIX B: METHODOLOGY	27
APPENDIX C: CODE SET DETAIL	30
APPENDIX D: POPULATION SAMPLE SIZE AND COST	36
APPENDIX E: NEW ONCOLOGY DRUG AND BIOLOGIC APPROVALS, 2004 – 2014	37
REFERENCES	40

EXECUTIVE SUMMARY

Nearly 14.5 million Americans with a history of cancer were alive in 2014¹ and that number is projected to increase to 18.1 million in 2020.² A number of factors will contribute to this increase, including the growth and aging of the U.S. population, an overall reduction in mortality, the earlier detection of cancer (lead time before death), and the increase in cancer survival. The projected increase for 2020 assumes continued past trends: the 5-year survival rate for all cancers diagnosed during 2005-2011 was 69%, up from 49% during 1975-1977,¹ and a 2012 study identified a 1.5% annual decline in cancer mortality for the decade examined.³

The increase in people living with cancer and the introduction of new therapies are associated with a rise in cancer care costs. Cancer care costs in the U.S. were estimated to be \$124.57 billion in 2010, and are projected to increase to \$158 to \$173 billion by 2020, representing a 27% to 39% increase.² In addition to the impact of increased cancer survivorship and new therapies, a significant shift in the site of chemotherapy infusion delivery from less expensive physician office settings to more expensive hospital outpatient settings has driven some of the past increase^{4,5,6,7,8} and may continue to contribute to cost increases.

The objective of this analysis was to identify trends in the overall and component costs of cancer care from 2004 to 2014 and to create comparisons to cost trends in the non-cancer population. We analyzed the prevalence and per-patient costs of actively treated cancer patients (those with claims for cancer surgery, radiation oncology, or chemotherapy) and non-actively treated cancer patients in each year from 2004 to 2014 using two data sources: the Medicare 5% sample claim database and the Truven MarketScan commercial claim database. Note that our analysis of the Medicare 5% sample claim database did not include pharmacy-dispensed drugs provided under the Part D benefit.

We identified the following key dynamics:

- 1. The percent increase in per-patient cost from 2004 to 2014 for actively treated Medicare fee-for-service (FFS) and commercially insured cancer patients has been similar to the corresponding increase for the respective non-cancer populations.
- 2. The per-patient cost of chemotherapy drugs is increasing at a much higher rate than other cost components of actively treated cancer patients, driven largely by biologics, but the chemotherapy drug increase has been offset by slower growth in other components.
- 3. The site of service for chemotherapy infusion has dramatically shifted from lower-cost physician office to higher-cost hospital outpatient settings.

We have important observations on trends in prevalence, cost, and site of service, summarized below:

 The prevalence of people living with cancer increased from 2004 to 2014 but the prevalence of patients receiving active treatment has remained relatively stable.

- Over the entire 2004 to 2014 study period, the average annual increase in cost was essentially the same in the actively treated cancer population and the non-cancer population.
- Cancer prevalence increased from 2004 to 2014 more than the contribution of cancer patients' cost to the total population spend.
- For patients being actively treated, the portion of spending for cancer-directed pharmaceuticals increased from 2004 to 2014 while the portion of spending for inpatient care declined.
 - In particular, the portion of spending for biologic chemotherapies increased from 3% to 9% in the Medicare population and from 2% to 7% in the commercial population.
- The portion of chemotherapy infusions being performed in generally more expensive hospital outpatient settings increased by at least 30%, from 2004 to 2014 with a corresponding reduction in the generally less expensive physician office settings.
- As explained in the body of the report, if the chemotherapy infusion site-of-service distribution in 2014 had been maintained at 2004 levels, the estimated Medicare FFS cost per infused chemotherapy patient in 2014 would have been approximately:
 - \$51,900 per actively treated Medicare FFS patient instead of the observed \$56,100 (7.5% lower)
 - \$89,900 per commercial patient instead of the \$95,400 observed (5.8% lower)

This analysis identifies several drivers influencing the rising cost of cancer care. We hope this material will encourage organizations to focus on feasible cost reduction opportunities.

This report was commissioned by Community Oncology Alliance, who received financial support from the following organizations: Bayer, Bristol-Myers Squibb, Eli Lilly and Company, Janssen Pharmaceuticals, Merck, Pfizer, Pharmaceutical Research and Manufacturers of America (PhRMA), and Takeda. The findings reflect the research of the authors; Milliman does not intend to endorse any product or organization. If this report is reproduced, we ask that it be reproduced in its entirety, as pieces taken out of context can be misleading. As with any economic or actuarial analysis, it is not possible to capture all factors that may be significant. Because we present national average data, the findings should be interpreted carefully before they are applied to any particular situation. These results are based on analysis of Truven MarketScan commercial data and the Medicare 5% sample data from 2004 to 2014. Different data sets, time periods, and methodologies will produce different results. Bruce Pyenson is a member of the American Academy of Actuaries and meets the Qualification Standards of the American Academy of Actuaries for this report.

4

BACKGROUND

The cost of cancer care has received significant attention in the last several years. Cancer care costs in the U.S. were estimated to be \$124.57 billion in 2010 and are projected to increase to \$158 to \$173 billion by 2020.² A study examining cancer care costs across 1987-2005 identified a near doubling of cancer care costs, but the study concluded this increase was not dissimilar to overall trends in aggregate health spending.⁹

The rise in the number of people living with a history of cancer from nearly 14.5 million in 2014¹ to an expected 18.1 million in 2020² will be a major driver of the increase in cancer care costs. The growth in cancer cases is driven in part by the growth and aging of the U.S. population while another significant driver has been the improvement in cancer survival. Improvements in cancer survival contribute to the rise in cancer prevalence rates: the 5-year survival rate for all cancers diagnosed during 2005-2011 was 69%, up from 49% during 1975-1977.¹ Earlier detection of cancer associated with improvements in cancer screening as well as innovations in cancer treatment are responsible for the improved survival rates. Even without improved survival, earlier detection can increase the number of people living with cancer due to what is called "lead time," increasing the time between cancer diagnosis and mortality. Cancer drug therapies have contributed to the rise in the cost of cancer care. Over 70 new drugs and biologics have been approved for cancer indications within the study period from 2004 to 2014 (a full listing is provided in Appendix E). In each of the past three years, more than 20 therapies have either been approved to treat cancer or received new cancer indications.¹⁰

Another trend contributing to the increase in cancer care costs has been the shift in the site of chemotherapy infusion delivery from generally lower-cost physician office settings to generally higher-cost hospital outpatient settings. Two site of care analyses using Truven MarketScan commercial data identified a 20% to 39% and a 28% to 53% lower cost per person receiving chemotherapy infusion in a physician office versus a hospital outpatient setting.^{7,8} An analysis of the Medicare FFS population identified a \$6,500 lower cost for Medicare beneficiaries receiving chemotherapy infusion in a physician office versus hospital outpatient setting.⁶

Along with this site of service shift, there has also been substantial consolidation among outpatient oncology providers and hospitals or health systems going back to at least 2003 with a notable increase starting in 2011. A recent study demonstrated that this increased provider consolidation resulted in statistically significant increased inflation-adjusted spending on outpatient prescription drug-based cancer treatment.¹¹ One recent study reported that hospitals participating in the federal 340B drug pricing program receive over 50 percent higher Part B oncology drug reimbursement per-beneficiary per-day than community oncology practices.⁵ The federal 340B drug pricing program allows hospitals to purchase drugs at greatly reduced prices; 340B was intended to support providers that furnish services to low-income people, allowing them to provide care to more patients using sometimes scarce federal resources.¹² However, providers can purchase drugs at these reduced prices for non-low-income patients, including those with Medicare or private insurance, and generate revenue if their reimbursement exceeds the price of the drug.¹²

Clinical progress is another potential driver of the cost of cancer care. For multiple myeloma, the 3-year survival was only 42% through the 1980s. In the 2000s, the introduction of thalidomide analogs and proteasome inhibitors increased that 3-year survival to approximately 66%, and

novel therapies and therapeutic mechanisms continue to be explored.¹³ Patients have also benefited from substantial progress in drug therapies treating colorectal cancer.¹⁴ These novel therapies may be associated with increased costs.

The objective of this analysis was to identify trends in cancer care costs and the components of cancer care costs from 2004 to 2014 while comparing those trends to that of the total population and non-cancer population. We analyzed actively treated cancer patients (those with claims for cancer surgery, radiation therapy, or chemotherapy) in each year from 2004 to 2014 in both the Medicare 5% sample and the Truven MarketScan commercial claim database.

FINDINGS

This section includes our detailed findings from 2004 to 2014 in the commercially insured and the Medicare FFS populations. There are three major sections to our findings:

- Cancer population characteristics
- o Spending on cancer
- Chemotherapy site of service

For more detail about our data sources and methodology please see the Appendices.

CANCER POPULATION CHARACTERISTICS: 2004-2014

Figure 1 splits the annual prevalence of cancer between actively treated patients and nonactively treated patients for 2004 to 2014.









Source: Based on Milliman analysis of the 2004-2014 Truven MarketScan data

The prevalence of cancer increased between 2004 and 2014 from 7.3% to 8.5% (a 16% increase) in the Medicare FFS population and from 0.7% to 0.9% (a 26% increase) in the commercially insured population. However, the prevalence of actively treated cancer remained relatively stable for both populations: 2.7% in 2004 and 2.6% in 2014 for the Medicare FFS population and 0.4% in both 2004 and 2014 for the commercial population.

^a Prevalence rates are measured as the number of patients with one or more specified claims coded with a cancer diagnosis in the given calendar year divided by the total database population. Actively treated cancer patients identified in each year include those with one or more claims for chemotherapy, radiation therapy or cancer surgery.

SPENDING ON CANCER

We examined the proportion of total spending contributed by the non-cancer, the actively treated cancer, and the non-actively treated cancer populations for each year of the study period. Table 1 presents the proportion each sub-population contributes to total population spend.

Table 1: Percent of total annual allowed costs by sub-population (actively treated cancer,
non-actively treated cancer, and non-cancer) ^b

Medicare FFS Population	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
Non-cancer	80.5%	80.7%	80.5%	80.5%	80.4%	80.6%	80.0%	79.4%	79.2%	79.3%	79.2%
Cancer	19.5%	19.3%	19.5%	19.5%	19.6%	19.4%	20.0%	20.6%	20.8%	20.7%	20.8%
Actively treated	11.6%	11.3%	11.3%	10.4%	10.9%	10.7%	11.0%	11.3%	11.4%	11.1%	11.2%
Non-actively treated	7.9%	7.9%	8.2%	9.1%	8.7%	8.7%	9.1%	9.4%	9.3%	9.6%	9.5%

Source: Based on Milliman analysis of the 2004-2014 Medicare 5% sample data

Commercially Insured Population	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
Non-cancer	90.6%	90.5%	90.3%	90.0%	89.7%	89.7%	89.4%	89.3%	89.2%	89.1%	89.3%
Cancer	9.4%	9.5%	9.7%	10.0%	10.3%	10.3%	10.6%	10.7%	10.8%	10.9%	10.7%
Actively treated	7.4%	7.3%	7.7%	7.9%	8.1%	8.1%	8.3%	8.4%	8.5%	8.5%	8.4%
Non-actively treated	2.0%	2.1%	2.0%	2.0%	2.2%	2.2%	2.3%	2.3%	2.3%	2.4%	2.3%

Source: Based on Milliman analysis of the 2004-2014 Truven MarketScan data

The percentage of total spending for cancer patients (both actively treated and non-actively treated) has increased slightly from 2004 to 2014 for both the Medicare and commercial populations. In the Medicare FFS population, the contribution to total population spend increased from 19.5% to 20.8% (6.7% increase), while in the commercial population, the contribution increased from 9.4% to 10.7% (13.8% increase).

Over the same period, the prevalence of cancer (actively treated and non-actively treated) increased at a higher rate than the increase in the spending contribution: prevalence increased from 7.3% to 8.5% (16% increase) in the Medicare population and from 0.7% to 0.9% (26% increase) in the commercially insured population (Figure 1). Cancer prevalence is increasing at a faster rate than the portion that the cancer population contributes to total population spend.

^b Allowed cost: all reimbursement from the payer plus member cost sharing. Note that for our Medicare analyses, pharmacy drugs covered under Part D are not included. For the commercial analysis, prescription drug costs are included and include oral chemotherapy drugs.

Actively treated cancer: members coded with cancer and having one or more claims for chemotherapy, radiation therapy or cancer surgery.

Non-actively treated cancer: members coded with cancer and not having one or more claims for chemotherapy, radiation therapy or cancer surgery. **Non cancer**: members without claims coded with cancer.

However, as shown in Figure 1, the increased prevalence reflects mostly an increase in patients who are not in active treatment.

Per-patient-per-year (PPPY) spending on actively treated cancer patients is higher than spending on non-actively treated patients and that spending is increasing (Figure 2).



Figure 2: PPPY allowed cost by subpopulation^c

^c **PPPY**: per patient per year.

Allowed cost: all reimbursement from the payer plus member cost sharing. Note that for our Medicare analyses, pharmacy drugs covered under Part D are not included. For the commercial analysis, prescription drug costs are included and include oral chemotherapy drugs. Total population: all members.

Non cancer: all members without claims coded with cancer.

Actively treated cancer: members coded with cancer and having one or more claims for chemotherapy, radiation therapy or cancer surgery. Non-actively treated cancer: members coded with cancer and not having one or more claims for chemotherapy, radiation therapy or cancer surgery.

Source: Based on Milliman analysis of the 2004-2014 Truven MarketScan data

While the higher PPPY cost for actively treated cancer patients is evident in Figure 2, the PPPY cost increased at a similar rate for all subpopulations. Figure 3 shows the cumulative trend in PPPY allowed cost for each subpopulation which is calculated as the trend from 2004 to each subsequent year.



Figure 3: Cumulative trend in PPPY allowed cost across subpopulations^d



2004-07

Actively treated cancer population

2004-08

d PPPY: per patient per year.

2004-05

Allowed cost: all reimbursement from the payer plus member cost sharing. Note that for our Medicare analyses, pharmacy drugs covered under Part D are not included. For the commercial analysis, prescription drug costs are included and include oral chemotherapy drugs.

2004-10

2004-11

Non-cancer population

2004-12

Non-actively treated cancer population

2004-13

2004-14

Total population: all members.

Non cancer: all members without claims coded with cancer.

2004-06

Total population

Actively treated cancer: members coded with cancer and having one or more claims for chemotherapy, radiation therapy or cancer surgery.

2004-09

Non-actively treated cancer: members coded with cancer and not having one or more claims for chemotherapy, radiation therapy or cancer surgery.

40%

30%

20%

10%

0%

Figure 3 demonstrates that per-patient costs for the total population, for the actively treated cancer population, and for the non-cancer population were increasing at very similar rates throughout the study period; 35.2% versus 36.4% and 34.8% respectively for the Medicare population and 62.9% versus 62.5% and 60.8% respectively for the commercial population. The non-actively treated cancer population's 10-year cost trend was noticeably lower for both the Medicare and commercial populations; 26.0% and 34.7% respectively. We calculated confidence intervals for each cohort's trend line using an exponential curve to fit the series of PPPYs. The three cohorts 95% confidence intervals overlap and by this measure the 10 year cost trend between the total population, non-cancer population and actively treated cancer population are not statistically different.

Components of cancer care

We split the PPPY cost for actively treated patients into cost categories to identify how specific components of cancer treatment have changed over the study period.

Figure 4 shows four pie charts, where per-patient annual allowed cost is split into eleven categories (described in Appendix B) for 2004 and 2014 for each of the Medicare and commercial populations.



Figure 4: PPPY allowed cost by cost category in the actively treated cancer population, Medicare and commercial[®]

Source: Based on Milliman analysis of the 2004-2014 Truven MarketScan data and Medicare 5% sample data

e PPPY: per patient per year.

Allowed cost: all reimbursement from the payer plus member cost sharing. Note that for our Medicare analyses, pharmacy drugs covered under Part D are not included. For the commercial analysis, prescription drug costs are included and include oral chemotherapy drugs. For a full explanation of the service categories used, see Appendix B.

In Figure 4, the bracketed pie sections shows the subtotal for chemotherapy drugs (biologic chemotherapy, cytotoxic chemotherapy, and other chemotherapy and cancer drugs). The portion of PPPY spending has increased in actively treated cancer populations, from 15% to 18% in Medicare and from 15% to 20% in commercial. In particular, biologic chemotherapies have seen an increase from 3% to 9% in the Medicare population and from 2% to 7% in the commercial population. The contribution of cost from hospital inpatient admissions decreased, from 27% to 24% for Medicare and from 21% to 18% for commercial. A similar decline was seen for contribution of cost from cancer surgeries, decreasing from 15% to 11% for Medicare and from 15% to 13% for commercial. Radiation oncology cost contribution for the Medicare population increase form 1% to 3%, but the portion did not increase for commercial.

Table 2 shows the change in PPPY allowed costs from 2004 to 2014 for the Medicare and commercial populations.

Table 2: 2004 to 2014 allowed cost trend by major service category for actively treated patients – Medicare and commercial^f

Service Category	2004-2014 PPP	Y Cost Trends
Service Calegory	Medicare	Commercial
Hospital Inpatient Admissions	22%	44%
Cancer Surgeries (inpatient and outpatient)	0%*	39%
Sub-Acute Services	51%	15%
Emergency Room	132%	147%
Radiology – Other	24%	77%
Radiation Oncology	204%	66%
Other Outpatient Services	48%	49%
Professional Services	40%	90%
Biologic Chemotherapy	335%	485%
Cytotoxic Chemotherapy	14%	101%
Other Chemo and Cancer Drugs	-9%	24%
Total PPPY Cost Trend	36%	62%

Source: Based on Milliman analysis of the 2004-2014 Truven MarketScan data and Medicare 5% sample data

The PPPY trends vary considerably by service category, with some categories (such as hospital inpatient admissions and other chemo and cancer drugs) increasing less than the total PPPY cost trend, and others (such as biologic chemotherapy) trending at much higher rates than the total PPPY cost trend. PPPY Medicare trends are lower for all services compared to commercial trends except sub-acute services and radiation oncology.

^fSee methodology Appendix B for definition of cost categories.

PPPY: per patient per year.

Allowed cost: all reimbursement from the payer plus member cost sharing. Note that for our Medicare analyses, pharmacy drugs covered under Part D are not included. For the commercial analysis, prescription drug costs are included and include oral chemotherapy drugs.

^{*} The difference in cost between 2004 and 2014 was \$6

Cost characteristics for specific cancer types

Table 3 shows the increase from 2004 to 2014 in annual per patient spending for specific types of cancer.

Table 3: 2004 to 2014 trend in PPPY allowed cost for actively treated cancer patients, by cancer type^g

Canaar Tura	2004-2014 PPP	Y Cost Trends
Cancer Type	Medicare	Commercial
Blood	53%	73%
Breast	36%	71%
Colon	28%	65%
Lung	21%	59%
Non-Hodgkin's Lymphoma	34%	69%
Pancreatic	25%	54%
Prostate	39%	79%
Other	22%	58%
Total: All Cancers	36%	62%

Source: Based on Milliman analysis of the 2004-2014 Truven MarketScan data and Medicare 5% sample data.

For the Medicare population, the 10-year trend in annual per-patient blood cancer and prostate cancer costs were higher than average, 53% and 39% respectively, while the 10-year trend in lung cancer, pancreatic cancer, and colon cancer costs were lower than average, 21%, 25% and 28% respectively. For the commercial population, the 10-year trend in colon cancer, blood cancer, breast cancer, and prostate cancer costs were higher than average, 65%, 73%, 71% and 79% respectively, while the 10-year trend in lung and pancreatic cancer costs were lower than average, 59% and 54% respectively.

Many factors can influence the change in costs, some of which involve shifts in treatment patterns for particular cancer types. The rise in costs for the blood cancer cohort during this period coincided with the introduction of FDA approved therapeutic options for patients with blood cancers such as myeloma and leukemia. In addition to improvements in survival, the use of new myeloma drugs led to a substantial increase in drug costs during this period.¹³

We examine colon cancer treatment costs in detail in Figure 5 to illustrate how therapeutic changes over the study period can influence costs. As seen in Table 3, the colon cancer population had lower than average cost trends for the Medicare population and higher than average cost trends for the commercial population. Figure 5 shows the components of annual per-patient cost for each year of the 11-year study period.

^g **PPPY**: per patient per year.

Allowed cost: all reimbursement from the payer plus member cost sharing. Note that for our Medicare analyses, pharmacy drugs covered under Part D are not included. For the commercial analysis, prescription drug costs are included and include oral chemotherapy drugs. See Appendix C for specific cancer identification criteria. Note that non-Hodgkin's lymphoma patients have been removed from the blood cancer category and are only included in the non-Hodgkin's lymphoma category.



Figure 5: 2004-2014 average allowed costs per actively treated colon cancer patient by claim category^h



Source: Based on Milliman analysis of the 2004-2014 Truven MarketScan data

h Allowed cost: all reimbursement from the payer plus member cost sharing. Note that for our Medicare analyses, pharmacy drugs covered under Part D are not included. For the commercial analysis, prescription drug costs are included and include oral chemotherapy drugs. See methodology Appendix B for colon cancer identification criteria and a description of claim categories.

For the actively treated colon cancer population, there was almost no spending on biologic therapies in 2004, but spending on these therapies increased dramatically in 2005. The following changes in treating colon cancer occurred over the study period:¹⁵

- In 2004, cetuximab was approved for the treatment of metastatic colon cancer.
- Panitumumab followed in 2006 for the treatment of epidermal growth factor receptor (EGFr)expressing metastatic colon cancer.
- In 2008, the ASCO annual meeting included five randomized controlled trials concluding that the presence of KRAS gene mutations in tumor tissue can predict the utility of these therapies, and the NCCN updated its guidelines in 2009.

CHEMOTHERAPY SITE OF SERVICE

Most payers, including Medicare, pay less if a chemotherapy infusion is provided in a physician office setting as opposed to a hospital outpatient facility setting. We examined changes from 2004 to 2014 in both spending and volume of chemotherapy infusion drugs by site of service. Figure 6 shows a large shift in spending on chemotherapy drugs to the generally more expensive hospital outpatient settings from generally less expensive physician office setting.



Figure 6: Infused chemotherapy drug spending by site of serviceⁱ

Source: Based on Milliman analysis of the 2004-2014 Truven MarketScan data and Medicare 5% sample data

As shown in Figure 7, from 2004 to 2014 hospital outpatient settings saw an increase in the portion of chemotherapy infusions, and physician offices saw a corresponding decrease in the portion of chemotherapy infusions. In Figure 7, we also include a 340B hospital setting category to capture the percentage of hospital outpatient department-based chemotherapy infusions that are being administered in hospitals participating in the 340B drug purchasing program. We provide 340B information for Medicare only as our commercial data does not have facility identifiers.

i PPPY: per patient per year.

Allowed cost: all reimbursement from the payer plus member cost sharing. Note that for our Medicare analyses, pharmacy drugs covered under Part D are not included. For the commercial analysis, prescription drug costs are included and include oral chemotherapy drugs. Other: site of service other than physician office and hospital outpatient includes SNF or home.







Source: Based on Milliman analysis of the 2004-2014 Medicare 5% sample data

Source: Based on Milliman analysis of the 2004-2014 Truven MarketScan data

The portion of chemotherapy infusions delivered in hospital outpatient departments increased from 15.8% to 45.9% in the Medicare population and 5.8% to 45.9% in the commercial population. In the Medicare population, the portion of chemotherapy infusions administered in a 340B hospital outpatient department increased from 3.0% to 23.1%. In 2014, chemotherapy infusions in 340B hospitals accounted for 50.3% of all chemotherapy infusions in hospital outpatient departments.

Figure 8 shows the PPPY allowed costs of chemotherapy patients by the site of where they received all of their chemotherapy infusions. For Medicare patients, this could be in a physician office only, a non-340B hospital outpatient department only, or a 340B hospital outpatient department only. For commercial patients, this could be in a physician office only or a hospital outpatient department department only. We excluded patients who received chemotherapy in both settings (~7%) and excluded pharmacy-based oral chemotherapy.



Figure 8: PPPY costs for chemotherapy patients based on site of their infused chemotherapy service, 2004 to 2014^j

Source: Based on Milliman analysis of the 2004-2014 Medicare 5% sample data



Source: Based on Milliman analysis of the 2004-2014 Truven MarketScan data

Figure 8 demonstrates that the average annual PPPY allowed cost for infused chemotherapy patients was significantly higher when chemotherapy infusions were delivered entirely in a hospital outpatient setting versus a physician office setting. Compared to patients receiving all chemotherapy infusions in a physician office, those receiving all chemotherapy infusions in a hospital outpatient facility had a PPPY that was \$13,167 (37%) higher in the 2004 Medicare

Allowed cost: all reimbursement from the payer plus member cost sharing. Note that for our Medicare analyses, pharmacy drugs covered under Part D are not included. For the commercial analysis, prescription drug costs are included and include oral chemotherapy drugs.

^j **PPPY**: per patient per year.

FFS population, \$16,208 (34%) higher in the 2014 Medicare FFS population, \$19,475 (25%) higher in the 2004 commercially insured population, and \$46,272 (42%) higher in the 2014 commercially insured population.

We note that the PPPYs for 340B hospitals and non-340B hospitals are similar. 340B hospitals receive the same payment for drugs from Medicare as non-340B hospitals but the purchase price from the drug manufacturer is at a mandated lower price than non-340B hospitals.

Modeling the cost implication of infused chemotherapy site of service shift

The shift to higher cost hospital outpatient departments has contributed to the rise in PPPY costs for both the Medicare and commercial populations. By applying the two factors involved - the site of service and the cost differential by site of service - we use a relatively simple approach to estimate the extra cost associated with the observed site of service shift.

While we believe this simple approach reasonably captures the extra cost associated with chemotherapy infusion in hospital outpatient settings versus physician office settings, we note the uncertainty in reducing total payer spending. In particular, if hospital outpatient departments had not expanded chemotherapy services, they may have found other ways to generate desired revenue.

We assumed the 2004 distribution of patients by the site of chemotherapy infusion would be maintained for each subsequent year. We applied the annual cost trends observed for each site of service in each year. This model captures the observed cost trends while maintaining the distribution of patients receiving chemotherapy in the two settings at 2004 levels. Figure 9 shows the observed versus the modeled PPPY costs over the study period.





Source: Based on Milliman analysis of the 2004-2014 Medicare 5% sample data



Source: Based on Milliman analysis of the 2004-2014 Truven MarketScan data

If the site of service distribution was the same in 2014 as in 2004, the average PPPY cost in 2014 for actively treated chemotherapy patients would have been about \$51,900 per Medicare beneficiary receiving infused chemotherapy instead of the \$56,000 observed (7.5% lower cost) and \$89,900 in the commercial population instead of the \$95,400 observed (5.8% lower cost).

k PPPY: per patient per year.

Allowed cost: all reimbursement from the payer plus member cost sharing. Note that for our Medicare analyses, pharmacy drugs covered under Part D are not included. For the commercial analysis, prescription drug costs are included and include oral chemotherapy drugs. Observed cost: PPPY cost of actively treated chemotherapy patients based on the actual distribution of chemotherapy infusion sites each year. Modeled cost: PPPY cost of actively treated chemotherapy patients if the 2004 distribution of chemotherapy infusion sites was maintained. For 2014, we estimate that Medicare spending would be about \$2 billion lower if the infused chemotherapy site of service shift had not occurred. Table 4 shows results of scenarios if varying portions of the observed shift had not occurred.

Table 4: Difference between observed and modeled Medicare FFS spending in 2014 for actively treated chemotherapy patients under site of service shifting scenarios

	Cost impact in billions in 2014				
	Shift to 25% of 2004 observed levels	Shift to 50% of 2004 observed levels	Shift to 75% of 2004 observed levels	Shift to 100% of 2004 observed levels	
Estimated Medicare FFS spending cost difference in 2014 if observed chemotherapy infusion site of service distribution was shifted toward 2004 site of service distribution	\$0.5	\$1.0	\$1.5	\$2.0	

Source: Based on Milliman analysis of the 2004-2014 Medicare 5% sample data. See Appendix D for Medicare population and cost.

In 2014, 64.6% of the Medicare actively treated cancer population was receiving infused chemotherapy. If the 2014 distribution of chemotherapy infusion site of service for these actively treated chemotherapy patients had been maintained at the 2004 levels for between 25% and 100% of these patients, the 2014 Medicare FFS spending would have been lower by between \$500 million and \$2 billion, respectively. Relative to 2014 Medicare spending, the \$2 billion lower cost represents:

- \circ 7.5% lower cost for the actively treated chemotherapy patients
- o 5.2% lower cost for the total actively treated cancer population
- o 0.59% lower cost for the total Medicare population spend

A number of factors could cause the modeled costs to be higher or lower than our estimate. We have assumed that the mix of patients receiving chemotherapy infusion in hospital outpatient facility settings and physician office settings were not substantially different. If the type of patients who shifted to receive their chemotherapy in a hospital outpatient facility setting were higher or lower need patients, the costs could vary from our modeled costs. However, differences in acuity for chemotherapy patients that shift from physician office to hospital outpatient to receive services is not substantiated in the literature. The modeled cost also does not consider regional variation in fee schedules for the commercial population or wage index, DSH or IME reimbursement differences for the Medicare population. If the shift occurred in regions with lower or higher reimbursement than the observed averages, the results could vary. The site of service estimated cost impact analysis does not account for the tendency for provider organizations to increase fees or utilization to meet revenue goals or the availability of alternate sites of service in all locales.

DISCUSSION AND CONCLUSIONS

Healthcare spending is in the headlines, including spending on cancer care. The overall increase in healthcare costs is widely viewed as unsustainable. Increased spending on healthcare by employers reduces funds available for wages; for government programs, increased spending reduces funds available for infrastructure or other investments. For cancer, prominent individuals associated with academic medicine have criticized the high price of particular therapies that they argue bring little clinical advantage to patients.

This paper explores very large administrative databases to identify the components of total annual spending on cancer care. Instead of focusing on the cost of individual patients or even cancer types, we look at the big picture of average annual per-patient spending on patients receiving cancer care. Our analysis, which focuses mainly on actively treated cancer patients and compares their trends over 11 years to the total population trends, demonstrates the following in particular:

1. The percent increase in cost from 2004-2014 for actively treated Medicare FFS and commercially insured cancer patients has been similar to the corresponding increase for the non-cancer Medicare FFS and commercially insured populations.

Our study finds that this increase is similar to the increase in overall population spending. While we observe that the prevalence of cancer has increased substantially in these populations over the study period, much of this increase is in the number of non-actively treated cancer patients.

2. The cost of chemotherapy drugs is increasing at a rate significantly higher than other cost components of actively treated cancer patients, driven largely by biologics, but the chemotherapy drug increase has been offset by slower growth in other component costs.

When cancer costs are separated into components to understand cost drivers, we see that chemotherapy drug costs are increasing more quickly than other components. These increasing drug costs have been offset by slower growth in other costs categories, such as inpatient services and cancer surgeries.

3. The site of service for chemotherapy infusion has dramatically shifted from physician office to hospital outpatient settings, which has contributed to the increase in cancer care cost.

The data demonstrates a consistent pattern of higher spending on patients receiving chemotherapy in hospital outpatient facilities than those receiving chemotherapy in physician office settings, and a trend toward the use of these higher cost hospital outpatient facility settings.

As healthcare spending continues to rise, insight into the drivers of cost in cancer care can enlighten strategies for meaningful change. Today's movement for systematic change is often referred to as population health and will certainly involve cancer care. We hope our big picture approach will be useful to that effort.

APPENDIX A: KEY DATA SOURCES

<u>Medicare 5% Sample.</u> The Medicare 5% sample is a limited data set containing all Medicare paid claims generated by a statistically-balanced sample of Medicare beneficiaries. Information includes county of residence, diagnosis codes, procedure codes, DRG codes, site of service information, beneficiary age, eligibility status and an indicator for HMO enrollment. Member identification codes are consistent from year to year and allow for multiyear longitudinal studies. The Medicare data does not include Part D prescription drug data. We used 2004-2014 data.

Truven Health Analytics MarketScan Commercial Claims Database. The Truven Health Analytics MarketScan Commercial Claims Database (MarketScan) contains all paid claims generated by 15-50 million commercially insured lives annually (depending on the year of data). The MarketScan database represents the inpatient and outpatient healthcare service use of individuals nationwide who are covered by the benefit plans of large employers, health plans, government, and public organizations. The data includes diagnosis codes, procedure codes, DRG codes, and NDC codes, along with site of service information and the amounts paid by commercial insurers. The MarketScan database links paid claims and encounter data to detailed patient information across sites and to types of providers. Patient identifiers are consistent over time, allowing for longitudinal studies. The annual medical database includes private sector health data from approximately 100 payers. We used the MarketScan data from 2004-2014.

APPENDIX B: METHODOLOGY

Steps for our claim data analysis included the following:

1. Identified all cancer patients in each year of analysis

Patients must have at least one acute inpatient or two observation, nonacute inpatient, emergency department, or outpatient claims on different days that contain a cancer ICD-9 diagnosis code in any position of the claim. See Appendix C for code sets.

2. Identified the subset of cancer patients being actively treated

Active treatment includes chemotherapy or radiation therapy or cancer surgery.

Patients were considered to have active chemotherapy treatment if they had 1+ claim for a chemotherapy J-code or chemotherapy NDC codes (commercial analysis only).

- For the chemotherapy J codes, we used the entire J8500 and J9000 series. The specific codes may have changed over the study period but the range has been maintained.
- Note that our analysis of the Medicare 5% sample claim database did not include pharmacy-dispensed drugs provided under the Part D benefit.

Patients were considered to have active inpatient chemotherapy treatment if they had a claim for any of the inpatient chemotherapy MS-DRGs.

Patients were considered to have active radiation oncology treatment if they have a claim for at least one of the radiation therapy codes included in Appendix C or for revenue code 333.

Patients were considered to have active surgical treatment if they met one of the following criteria:

- Inpatient: Surgical MS-DRG that is coded with a Cancer ICD-9 code in the primary position of the claim
- Outpatient: Outpatient cancer surgery coded with a cancer procedure code and coded with a cancer ICD-9 code in the primary diagnosis of the claim

3. Identified characteristics of the actively treated cancer patients in each annual cohort

Major cancer types

- Specific cancers identified as cancer cases with at least two claims coded with the relevant ICD-9 diagnosis code for that cancer type in the primary position of the claim.
 For patients coded with more than one cancer type (approximately 1% of patients), we applied the following hierarchy:
 - Lung
 - Pancreatic
 - Blood
 - Non-Hodgkin's lymphoma

- Colon
- Breast
- Prostate
- Other

Chemotherapy site of service

- Percent of chemotherapy rendered in a hospital outpatient versus physician office setting
 - Place of service code 22 represents hospital outpatient departments, and 11 represents physician office
- Percent of hospital outpatient chemotherapy delivered in 340B hospitals
 - Note: the source for information about 340B status is the covered entities file, which is available for download at opanet.hrsa.gov and contains quarter-byquarter 340B eligibility for all acute care hospitals in the country.

4. Characterized costs by major service categories

All annual claims for each actively treated cancer patient were grouped into cost model categories based on ICD-9 procedure codes, HCPCs codes, CPT codes, revenue codes, place of service codes, and DRGs.

Categories Used in Exhibits	Description
Biologic chemotherapy drugs	Includes all biologic chemotherapy drugs; to identify, see infused chemotherapy codes and chemotherapy NDC codes lists; type of chemotherapy identified
Cancer surgeries (inpatient and outpatient)	Includes all surgical admissions (with a surgical DRG or MS-DRG) to inpatient acute care hospitals with a cancer diagnosis code as the primary ICD-9 diagnosis code and outpatient surgeries as identified on the outpatient cancer surgeries code list; must also have cancer ICD-9 diagnosis code as primary
Cytotoxic chemotherapy drugs	Includes all cytotoxic chemotherapy drugs; to identify, see infused chemotherapy codes and chemotherapy NDC codes lists; type of chemotherapy identified
Emergency room	Includes standalone emergency room visits (not resulting in an inpatient admission)
Hospital inpatient admissions	Includes all admissions to inpatient acute care hospitals with the exception of cancer surgeries, including medical admissions, non-cancer surgical admissions, rehabilitation and psychiatric admissions, and all associated professional services
Other chemo and cancer- related drugs	Includes all hormonal or other chemotherapy drugs (to identify, see infused chemotherapy codes and chemotherapy NDC codes lists; type of chemotherapy identified) as well as chemotherapy adjuncts (see

Cost model categories include:

	chemotherapy adjuncts code list) and hematopoietic agents (see hematopoietic agent J-code list and hematopoietic agent NDC code list)
Other outpatient services	Includes all laboratory and pathology services and outpatient procedural care (such as port placement or non-chemotherapy infusion services), as well as any non-cancer drugs and administration services. Includes non- chemotherapy prescription drug costs for commercial population
Professional services (office visits, urgent care, and chemotherapy administration)	Includes all professional E/M charges, whether in office or hospital outpatient setting, as well as all chemotherapy administration services as identified chemotherapy administration list
Radiation oncology See radiation oncology code list	
Radiology - other All radiology services, including those on the high tech imaging code list	
Sub-acute services (home health, hospice, and SNF)Includes all skilled nursing facility services, home health services, a hospice services	

APPENDIX C: CODE SET DETAIL

The following CPT and Revenue codes were used to identify claims as Acute Inpatient, Observation, Nonacute Inpatient, Emergency Department, or Outpatient site of service.

Claim type	CPT code	Revenue codes
Outpatient	99201-99205, 99211-99215, 99241-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99420, 99429, 99455, 99456, G0402, 0438,G0439,G0463,T1015	0510-0517,0519-0523, 0526-0529, 0982, 0983
Non-acute inpatient	99304-99310, 99315, 99316, 99318, 99324- 99328, 99334-99337	0118, 0128, 0138, 0148, 0158, 0190- 0194,0199, 0524, 0525, 0550-0552, 0559,0660-0663,0669
Acute inpatient	99221-99223, 99231-99233, 99238, 99239, 99251-99255, 99291, 99468, 99469, 99471, 99472, 99475-99480	010x, 0110-0115, 0117, 0119-0125, 0127, 0129-0135, 0137,0139-0145, 0147, 0149- 0155, 0157,0159-0162, 0164, 0166-0175, 0179, 0200-0204, 0206-0214, 0219, 0720- 0724, 0729, 0987
Observation 99217-99220		
Emergency department	99281-99285	0450-0452, 0456, 0459, 0981

Cancer ICD-9 codes

ICD-9 Diagnosis Code	Descriptor
140.xx-172.xx	Primary malignant neoplasms, not lymphatic or hematopoietic
174.xx-195.xx	Primary malignant neoplasms, not lymphatic or hematopoietic
196.xx-198.xx	Secondary malignant neoplasms (i.e., metastatic)
199.xx	Malignant neoplasms, unknown site
200.xx-208.xx	Leukemias and lymphomas
209.0x-209.3x	Neuroendocrine tumors
230.xx-234.xx	Carcinoma in situ

Specific cancer type ICD-9 codes

ICD-9 Diagnosis Code	Descriptor
162.xx	Lung Cancer
174.xx, 233.0	Breast Cancer
185.xx	Prostate Cancer
157.xx	Pancreatic Cancer
153.xx	Colon Cancer
200.xx, 202.0x-202.2x, 202.7x-202.8x	Non-Hodgkin Lymphoma
202.4x, 203.1x, 204.xx-208.xx	Blood Cancer

Chemotherapy DRGs and MS-DRGs

For discharges on or after 10/1/2007, used the MS-DRG list. For discharges before 10/1/2007, used the DRG list.

MS-DRG	MS-DRG Title
837	CHEMO W ACUTE LEUKEMIA AS SDX OR W HIGH DOSE CHEMO AGENT W MCC
838	CHEMO W ACUTE LEUKEMIA AS SDX W CC OR HIGH DOSE CHEMO AGENT
839	CHEMO W ACUTE LEUKEMIA AS SDX W/O CC/MCC
846	CHEMOTHERAPY W/O ACUTE LEUKEMIA AS SECONDARY DIAGNOSIS W MCC
847	CHEMOTHERAPY W/O ACUTE LEUKEMIA AS SECONDARY DIAGNOSIS W CC
848	CHEMOTHERAPY W/O ACUTE LEUKEMIA AS SECONDARY DIAGNOSIS W/O CC/MCC
DRG	DRG Title
410	CHEMOTHERAPY W/O ACUTE LEUKEMIA AS SECONDARY DIAGNOSIS
492	CHEMOTHERAPY W ACUTE LEUKEMIA OR W USE OF HI DOSE CHEMOAGENT

Radiation therapy codes

CPT code range	Description
77261-77263	Therapeutic Radiology: Treatment Planning
77280-77299	Radiation Therapy Simulation
77300-77370	Radiation Physics Services
77371-77373	Stereotactic Radiosurgery (SRS) Planning and Delivery
77399	Unlisted procedure, medical radiation physics, dosimetry and treatment devices,
11399	and special services
77401-77417	Radiation Treatment
77418	IMRT Delivery
77421	Stereoscopic Imaging Guidance
77422-77423	Neutron Therapy
77427-77499	Radiation Therapy Management
77520-77525	Proton Therapy
77600-77620	Hyperthermia Treatment
77750-77799	Brachytherapy

Chemotherapy J codes

HCPCS	Description	Category
J8510	Oral busulfan	Cytotoxic
J8520	Capecitabine, oral, 150 mg	Cytotoxic
J8521	Capecitabine, oral, 500 mg	Cytotoxic
J8530	Cyclophosphamide oral 25 MG	Cytotoxic
J8560	Etoposide oral 50 MG	Cytotoxic
J8561	Oral everolimus	Cytotoxic
J8562	Oral fludarabine phosphate	Cytotoxic
J8565	Gefitinib oral	Cytotoxic
J8600	Melphalan oral 2 MG	Cytotoxic
J8610	Methotrexate oral 2.5 MG	Cytotoxic
J8700	Temozolomide	Cytotoxic
J8705	Topotecan oral	Cytotoxic
J8999	Oral prescription drug chemo	Cytotoxic

10000	Develophing that in its other	Outotavia
J9000	Doxorubicin hcl injection	Cytotoxic
J9001	Doxorubicin hcl liposome inj	Cytotoxic
J9002	Doxil injection	Cytotoxic
J9010	Alemtuzumab injection	Biologic
J9015	Aldesleukin injection	Biologic
J9017	Arsenic trioxide injection	Cytotoxic
J9019	Erwinaze injection	Biologic
J9020	Asparaginase, NOS	Biologic
J9025	Azacitidine injection	Cytotoxic
J9027	Clofarabine injection	Cytotoxic
J9031	Bcg live intravesical vac	Biologic
J9033	Bendamustine injection	Cytotoxic
J9035	Bevacizumab injection	Biologic
J9040	Bleomycin sulfate injection	Cytotoxic
J9041	Bortezomib injection	Cytotoxic
J9042	Brentuximab vedotin inj	Biologic
J9043	Cabazitaxel injection	Cytotoxic
J9045	Carboplatin injection	Cytotoxic
J9047	Injection, carfilzomib, 1 mg	Cytotoxic
J9050	Carmustine injection	Cytotoxic
J9055	Cetuximab injection	Biologic
J9060	Cisplatin 10 MG injection	Cytotoxic
J9062	Cisplatin 50 MG injection	Cytotoxic
J9065	Inj cladribine per 1 MG	Cytotoxic
J9070	Cyclophosphamide 100 MG inj	Cytotoxic
J9080	Cyclophosphamide 200 MG inj	Cytotoxic
J9090	Cyclophosphamide 500 MG inj	Cytotoxic
J9091	Cyclophosphamide 1.0 grm inj	Cytotoxic
J9092	Cyclophosphamide 2.0 grm inj	Cytotoxic
J9093	Cyclophosphamide lyophilized	Cytotoxic
J9094	Cyclophosphamide lyophilized	Cytotoxic
J9095	Cyclophosphamide lyophilized	Cytotoxic
J9096	Cyclophosphamide lyophilized	Cytotoxic
J9097	Cyclophosphamide lyophilized	Cytotoxic
J9098	Cytarabine liposome inj	Cytotoxic
J9100	Cytarabine hcl 100 MG inj	Cytotoxic
J9110	Cytarabine hcl 500 MG inj	Cytotoxic
J9120	Dactinomycin injection	Cytotoxic
J9130	Dacarbazine 100 mg inj	Cytotoxic
J9140	Dacarbazine 200 MG inj	Cytotoxic
J9150	Daunorubicin injection	Cytotoxic
J9151	Daunorubicin citrate inj	Cytotoxic
J9155	Degarelix injection	Hormonal
J9160	Denileukin diftitox inj	Biologic
J9165	Diethylstilbestrol injection	Hormonal
J9103	Docetaxel injection	Cytotoxic
J9170 J9171	Docetaxel injection	Cytotoxic
J9171	Elliotts b solution per ml	Not
J9178	Inj, epirubicin hcl, 2 mg	Cytotoxic
J9179	Eribulin mesylate injection	Cytotoxic
J9181	Etoposide injection	Cytotoxic
J9182	Etoposide injection	Cytotoxic
J9185	Fludarabine phosphate inj	Cytotoxic

104.00		
J9190	Fluorouracil injection	Cytotoxic
J9200	Floxuridine injection	Cytotoxic
J9201	Gemcitabine hcl injection	Cytotoxic
J9202	Goserelin acetate implant	Hormonal
J9206	Irinotecan injection	Cytotoxic
J9207	Ixabepilone injection	Cytotoxic
J9208	Ifosfamide injection	Cytotoxic
J9209	Mesna injection	Not
J9211	Idarubicin hcl injection	Cytotoxic
J9212	Interferon alfacon-1 inj	Biologic
J9213	Interferon alfa-2a inj	Biologic
J9214	Interferon alfa-2b inj	Biologic
J9215	Interferon alfa-n3 inj	Biologic
J9216	Interferon gamma 1-b inj	Biologic
J9217	Leuprolide acetate suspension	Hormonal
J9218	Leuprolide acetate injection	Hormonal
J9219	Leuprolide acetate implant	Hormonal
J9225	Vantas implant	Hormonal
J9226	Supprelin LA implant	Hormonal
J9228	Ipilimumab injection	Biologic
J9230	Mechlorethamine hcl inj	Cytotoxic
J9245	Inj melphalan hydrochl 50 MG	Cytotoxic
J9250	Methotrexate sodium inj	Cytotoxic
J9260	Methotrexate sodium inj	Cytotoxic
J9261	Nelarabine injection	Cytotoxic
J9262	Inj, omacetaxine mep, 0.01mg	Cytotoxic
J9263	Oxaliplatin	Cytotoxic
J9264	Paclitaxel protein bound	Cytotoxic
J9265	Paclitaxel injection	Cytotoxic
J9266	Pegaspargase injection	Biologic
J9268	Pentostatin injection	Cytotoxic
J9270	Plicamycin (mithramycin) inj	Cytotoxic
J9280	Mitomycin injection	Cytotoxic
J9290	Mitomycin 20 MG inj	Cytotoxic
J9291	Mitomycin 40 MG inj	Cytotoxic
J9293	Mitoxantrone hydrochl / 5 MG	Cytotoxic
J9300	Gemtuzumab ozogamicin inj	Biologic
J9302	Ofatumumab injection	Biologic
J9303	Panitumumab injection	Biologic
J9305	Pemetrexed injection	Cytotoxic
J9306	Injection, pertuzumab, 1 mg	Biologic
J9307	Pralatrexate injection	Cytotoxic
J9310	Rituximab injection	Biologic
J9315	Romidepsin injection	Cytotoxic
J9320	Streptozocin injection	Cytotoxic
J9328	Temozolomide injection	Cytotoxic
J9330	Temsirolimus injection	Cytotoxic
J9340	Thiotepa injection	Cytotoxic
J9350	Topotecan injection	Cytotoxic
J9351	Topotecan injection	Cytotoxic
J9354	Inj, ado-trastuzumab emt 1mg	Biologic
J9355	Trastuzumab injection	Biologic
	Valrubicin injection	Cytotoxic
J9357		Cyluluxic

J9360	Vinblastine sulfate inj	Cytotoxic
J9370	Vincristine sulfate 1 MG inj	Cytotoxic
J9371	Inj, vincristine sul lip 1mg	Cytotoxic
J9375	Vincristine sulfate 2 MG inj	Cytotoxic
J9380	Vincristine sulfate 5 MG inj	Cytotoxic
J9390	Vinorelbine tartrate inj	Cytotoxic
J9395	Injection, Fulvestrant	Hormonal
J9400	Inj, ziv-aflibercept, 1mg	Biologic
J9600	Porfimer sodium injection	Other
J9999	Chemotherapy drug	Cytotoxic
Q2043	Provenge, 50 million autologous CD54+ cells	Biologic
Q2049	Lipodox 10 mg	Cytotoxic
Q2050	Doxil 10mg	Cytotoxic
Q0138	Ferumoxytol	Not

Outpatient cancer surgeries

Procedure code	Code description
11600	Exc tr-ext mal+marg 0.5 cm/<
11601	Exc tr-ext mal+marg 0.6-1 cm
11602	Exc tr-ext mal+marg 1.1-2 cm
11603	Exc tr-ext mal+marg 2.1-3 cm
11604	Exc tr-ext mal+marg 3.1-4 cm
11606	Exc tr-ext mal+marg >4 cm
11620	Exc h-f-nk-sp mal+marg 0.5/<
11621	Exc s/n/h/f/g mal+mrg 0.6-1
11622	Exc s/n/h/f/g mal+mrg 1.1-2
11623	Exc s/n/h/f/g mal+mrg 2.1-3
11624	Exc s/n/h/f/g mal+mrg 3.1-4
11626	Exc s/n/h/f/g mal+mrg >4 cm
11640	Exc f/e/e/n/l mal+mrg 0.5cm<
11641	Exc f/e/e/n/l mal+mrg 0.6-1
11642	Exc f/e/e/n/l mal+mrg 1.1-2
11643	Exc f/e/e/n/l mal+mrg 2.1-3
11644	Exc f/e/e/n/l mal+mrg 3.1-4
11646	Exc f/e/e/n/l mal+mrg >4 cm
17304	Mohs 1 stage
17305	Mohs 2 stage
17306	Mohs 3 stage
17307	Mohs addl stage
17310	Mohs addl specimen
17311	Mohs 1 stage h/n/hf/g
17312	Mohs addl stage
17313	Mohs 1 stage t/a/l
17314	Mohs addl stage t/a/l
17315	Mohs surg addl block
19160	Partial mastectomy
19162	P-mastectomy w/ln removal
19200	Mast radical
19220	Mast rad urban type

400.40	
19240	Mast mod rad
19301	Partial mastectomy
19302	P-mastectomy w/In removal
19305	Mast radical
19306	Mast rad urban type
19307	Mast mod rad
45384	Colonoscopy w/lesion removal
45385	Colonoscopy w/lesion removal
44139	Mobilization of colon
44140	Partial removal of colon
44141	Partial removal of colon
44143	Partial removal of colon
44144	Partial removal of colon
44145	Partial removal of colon
44146	Partial removal of colon
44147	Partial removal of colon
44150	Removal of colon
44151	Removal of colon/ileostomy
44152	Colectomy w/ileoanal anast
44153	Colectomy w/ileoanal anast
44155	Removal of colon/ileostomy
44156	Removal of colon/ileostomy
44157	Colectomy w/ileoanal anast
44158	Colectomy w/neo-rectum pouch
44160	Removal of colon
44204	Laparo partial colectomy
44205	Lap colectomy part w/ileum
44206	Lap part colectomy w/stoma
44207	L colectomy/coloproctostomy
44208	L colectomy/coloproctostomy
44210	Laparo total proctocolectomy
44211	Lap colectomy w/proctectomy
44212	Laparo total proctocolectomy
44213	Lap mobil splenic fl add-on
58150	Total hysterectomy
58152	Total hysterectomy
58180	Partial hysterectomy
58200	Extensive hysterectomy
58210	Extensive hysterectomy
58240	Removal of pelvis contents
58260	Vaginal hysterectomy
58262	Vag hyst including t/o
58263	Vag hyst w/t/o & vag repair
58267	
	Vag hyst w/urinary repair
58270	Vag hyst w/enterocele repair
58275	Hysterectomy/revise vagina
58280	Hysterectomy/revise vagina
58285	Extensive hysterectomy
58290	Vag hyst complex
58291	Vag hyst incl t/o complex
58292	Vag hyst t/o & repair compl
58293	Vag hyst w/uro repair compl
58294	Vag hyst w/enterocele compl

APPENDIX D: POPULATION SAMPLE SIZE AND COST

Demosmenteire	Com	nercial	Medi	care*
Demographics	2004	2014	2004	2014
Total population (number of members)	9,365,890	30,736,563	1,614,417	1,566,804
Cancer population				
Number of members	63,935	264,204	118,089	133,225
Percent of total population	0.7%	0.9%	7.3%	8.5%
Actively treated cancer population				
Number of members	34,329	129,507	44,139	41,098
Percent of total population	0.4%	0.4%	2.7%	2.6%
Percent of cancer population	53.7%	49.0%	37.4%	30.8%
Total chemotherapy population				
Number of members	24,968	102,130	27,556	26,563
Percent of actively treated cancer population	72.7%	78.9%	62.4%	64.6%
Total infused chemotherapy population				
Infusions in physician's office only				
Number of members	10,778	23,708	22,295	14,854
Percent of total chemotherapy population	43.2%	23.2%	80.9%	55.9%
Infusions in hospital outpatient facility only				
Number of members	817	20,258	3,622	10,047
Percent of total chemotherapy population	3.3%	19.8%	13.1%	37.8%
Infusions in a combination of settings				
Number of members	1,773	7,525	1,639	1,662
Percent of total chemotherapy population	7.1%	7.4%	5.9%	6.3%
Spending				
Total population (allowed cost)	\$24,379,270,607	\$129,785,018,754	\$13,446,397,168	\$17,670,384,548
Cancer population				
Total spending (allowed)	\$2,281,981,711	\$13,908,337,950	\$2,619,153,436	\$3,672,799,298
Percent of total population spend	9.4%	10.7%	19.5%	20.8%
Actively treated cancer population	-	1		1
Total spending (allowed)	\$1,794,163,969	\$10,915,304,244	\$1,554,903,891	\$1,986,161,660
Percent of total population spend	7.4%	8.4%	11.6%	11.2%
Percent of cancer population spend	78.6%	78.5%	59.4%	54.1%
Total chemotherapy population				
Total spending (allowed)	\$1,381,572,624	\$9,052,690,542	\$1,020,380,061	\$1,397,000,065
Percent of total population spend	5.7%	7.0%	7.6%	7.9%
Percent of cancer population spend	60.5%	65.1%	39.0%	38.0%
Percent of actively treated population spend	77.0%	82.9%	65.6%	70.3%
Total infused chemotherapy population	-			
Physician's office only				
Total spending (allowed)	\$758,511,113	\$2,361,766,202	\$753,370,260	\$668,629,519
Spending as % of total pop. spend	3.1%	1.8%	5.6%	3.8%
Spending as % of cancer pop. spend	33.2%	17.0%	28.8%	18.2%
Spending as % of actively treated pop. spend	42.3%	21.6%	48.5%	33.7%
Spending as % chemotherapy pop. spend	54.9%	26.1%	73.8%	47.9%
Hospital outpatient facility only				
Total spending (allowed)	\$72,796,282	\$2,843,556,239	\$162,497,323	\$595,661,897
Spending as % of total pop. spend	0.3%	2.2%	1.2%	3.4%
Spending as % of cancer pop. spend	3.2%	20.4%	6.2%	16.2%
Spending as % of actively treated pop. spend	4.1%	26.1%	10.5%	30.0%

Source: Based on Milliman analysis of the 2004-2014 Truven MarketScan data and Medicare 5% sample data

APPENDIX E: NEW ONCOLOGY DRUG AND BIOLOGIC APPROVALS, 2004 – 2014

The table below lists all drugs approved with cancer indications between 2004 and 2014, as extracted from the *New Drug Approvals* documentation prepared annually by the Pharmaceutical Research and Manufacturers of America.

Name (active ingredient)	Indication	FDA approval date
Alimta (pemetrexed)	Treatment of malignant pleural mesothelioma (asbestos-related cancer)	2/4/2004
Avastin (bevacizumab)	Treatment of first-line or previously untreated metastatic colorectal cancer Treatment of children with refractory or relapsed acute lymphoblastic	2/26/2004
Clolar (clofarabine)	leukemia	12/28/2004
Erbitux (cetuximab)	Treatment of metastatic colorectal cancer Stimulation of pancreatic secretions to aid in the diagnosis of pancreatic exocrine dysfunction, stimulation of gastrin secretion to aid in the	2/12/2004
Human Secretin (for injection)	diagnosis of gastrinoma, and stimulation of pancreatic secretion to facilitate the identification of the ampulla of Vater and accessory papilla during endoscopic retrograde cholangiopancreatography	4/9/2004
Sensipar (cinacalcet HCI)	Treatment of secondary hyperthyroidism in chronic kidney disease in patients on dialysis and for treatment of hypercalcemia in patients with parathyroid carcinoma	3/8/2004
Tarceva (erlotinib)	Treatment of advanced or metastatic non-small-cell lung cancer (NSCLC)	11/18/2004
Arranon (nelarabine)	Treatment of T-cell acute lymphoblastic leukemia and T-cell lymphoblastic lymphoma	10/28/2005
Nexavar (sorafenib)	Treatment of advanced renal cell carcinoma (kidney cancer)	12/20/2005
Sprycel (dasatinib)	Treatment of chronic myeloid leukemia and Philadelphia chromosome- positive acute lymphoblastic leukemia	6/28/2006
Sutent (sunitinib)	Treatment of advanced kidney cancer and gastrointestinal stromal tumors (GIST)	1/26/2006
Vectibix (panitumumab)	Treatment of epidermal growth factor receptor (EGFr)-expressing metastatic colorectal cancer	9/27/2006
Zolinza (vorinostate) Gardasil (quadrivalent human	Treatment of cutaneous manifestations in cutaneous T-cell lymphoma	10/6/2006
papillomavirus types 6,11,16,	Vaccination in females 9-26 years of age for prevention of certain	6/8/2006
and 18 vaccine) Ixempra (iabepilone)	diseases caused by human papillomavirus (HPV) types 6, 11, 16, and 18 monotherapy of metastatic or locally advanced breast cancer	10/16/2007
Tasigna (nilotinib)	Treatment of Philadelphia chromosome-positive chronic myeloid leukemia	10/29/2007
Torisel (temsirolimus)	Treatment of advanced renal cell carcinoma	5/30/2007
Tykerb (lapatinib)	Once-daily treatment of advanced or metastatic breast cancer in combination with Xeloda®	3/13/2007
Degarelix (for injection)	Treatment of advanced prostate cancer	12/24/2008
Mozobil (plerixafor)	To mobilize stem cells for autologous transplantation in non-Hodgkin's	12/15/2008
Treanda (bendamustine)	lymphoma and multiple myeloma Treatment of chronic lymphocytic leukemia	3/20/2008
Afinitor (everolimus)	Treatment of advanced renal cell carcinoma	3/30/2009
Arzerra (ofatumumab)	Treatment of refractory chronic lymphocytic leukemia (CLL)	10/26/2009
Folotyn (pralatrexate)	Treatment of relapsed or refractory peripheral T-cell lymphoma	9/24/2009
Istodax (romidepsin)	Treatment of cutaneous T-cell lymphoma (CTCL)	11/5/2009
Votrient (pazopanib)	Treatment of advanced renal cell carcinoma	10/19/2009
Halaven (eribulin mesylate injection)	Treatment of late-stage metastatic breast cancer	11/15/2010
Jevtana (cabazitaxel injection)	Treatment of metastatic hormone-refractory prostate cancer	6/17/2010
Provenge (sipuleucel-T)	Treatment of metastatic hormone-refractory prostate cancer	4/29/2010
Adcetris (brentuximab vedotin) Caprelsa (vandetanib tablets)	Hodgkin lymphoma, systemic anaplastic large-cell lymphoma Medullary thyroid cancer	8/19/2011 4/6/2011

Erwinaze (asparginase)	Acute lymphoblastic leukemia (ALL)	11/18/2011
Jakafi (ruxolitinib tablets)	myelofibrosis	11/16/2011
Xalkori (crizotinib)	Non-small-cell lung cancer	8/26/2011
Yervoy (ipilimumab injection for intravenous infusion)	Metastatic melanoma	3/25/2011
Zelboraf (vemurafenib tablets)	Metastatic melanoma	8/17/2011
Zytiga (abiraterone acetate		4/28/2011
tablets)	Metastatic prostate cancer	1/20/2011
Bosulif (bosutinib)	Treatment of previously treated Philadelphia chromosome-positive (Ph+) chronic myelogenous leukemia (CML) (chronic, accelerated or blast	9/4/2012
	phase)	5/4/2012
Choline C 11 injection	PET imaging for detection of recurrent	9/12/2012
	prostate cancer	5/12/2012
Cometriq (cabozantinib)	Treatment of metastatic medullary thyroid cancer	11/29/2012
Erivedge (vismodegib)	Treatment of advanced basal cell carcinoma	1/30/2012
Iclusig (ponatinib)	Treatment of CML and Philadelphia chromosome-positive acute	12/14/2012
	lymphoblastic leukemia (Ph+ALL)	
Inlyta (axitinib) Kyprolis (carfilzomib)	Treatment of advanced renal cell carcinoma Treatment of multiple myeloma	1/27/2012 7/20/2012
	Treatment of HER2-positive metastatic	
Perjeta (pertuzumab)	breast cancer	6/8/2012
Picato (ingenol mebutate)	Treatment of actinic keratosis	1/23/2012
Stivarga (regorafenib) Synribo (omacetaxine	Treatment of advanced colorectal cancer	9/27/2012
mepesuccinate)	Treatment of CML (chronic or accelerated phase)	10/26/2012
Tbo-filgrastim	Treatment of chemotherapy-induced neutropenia	8/29/2012
Voraxaze (glucarpidase)	Treatment of toxic levels of methotrexate in	1/17/2012
(g)	blood due to kidney failure Treatment of advanced castration-resistant	
Xtandi (enzalutamide)	prostate cancer	8/31/2012
Zaltrap (ziv-aflibercept)	Treatment of previously treated metastatic	8/3/2012
	colorectal cancer	
Gazyva (obinutuzumab) Gilotrif (afatinib)	Chronic lymphocytic leukemia EGFR-positive non-small-cell lung cancer	11/1/2013 7/12/2013
Imbruvica (ibrutinib)	Mantle cell lymphoma	11/13/2013
Kadcyla (ado-trastuzumab		2/22/2013
emtansine)	HER2-positive metastatic breast cancer	2/22/2013
Lymphoseek (technetium Tc99m tilmanocept)	lymphatic mapping in breast cancer and melanoma patients	3/13/2013
Mekinist (trametinib)	Unresectable or metastatic melanoma	5/29/2013
Pomalyst (pomalidomide)	Relapsed and refractory multiple myeloma	2/8/2013
Tafinlar (dabrafenib)	Unresectable or metastatic melanoma	5/29/2013
Xofigo (radium Ra 223 dichloride)	Castration resistant prostate cancer	5/15/2013
Beleodaq (belinostst)	Castration-resistant prostate cancer Treatment of relapsed or refractory peripheral T-cell lymphoma (PTCL)	7/3/2014
Blincyto (blinatumomab)	Treatment of Philadelphia chromosome-negative relapsed or refractory	12/3/2014
Bincyto (binatumomab)	B-cell precursor acute lymphoblastic leukemia (Ph-ALL)	12/3/2014
Cyramza (ramucirumab)	Treatment of advanced gastric cancer or gastroesophageal junction adenocarcinoma	4/21/2014
Keytruda (pembrolizumab)	Treatment of unresectable or metastatic melanoma (second-line therapy)	9/4/2014
Lynparza (olaparib)	Treatement of advanced ovarian cancer in patients with germline BRCA-	12/19/2014
	mutations	
Opdivo (nivolumab)	Treatment of unresectable or metastatic melanoma (second-line therapy) Treatment of relapsed follicular B-cell non-Hodgkin lymphoma, relapsed	12/22/2014
Zydelig (idelalisib)	small lymphocytic lymphoma	7/23/2014
Zydelig (idelalisib)	Treatment of relapsed chronic lymphocytic leukemia	7/23/2014

Zykadia (ceritinib)	Treatment of metastatic anaplastic lymphoma kinase-positive (ALK+) metastatic non-small cell lung cancer (NSCLC)	4/29/2014
Gardasil ®9 (human papillomavirus 9-valent vaccine, recombinant)	For the prevention of cervical vulvar, vaginal and anal cancers caused by nine types of human papillomavirus (HPV)	12/10/2014

REFERENCES

- 1. The American Cancer Society. Cancer Facts & Figures 2016. Atlanta: The American Cancer Society; 2016.
- Mariotto AB, Yabroff KR, Shao Y, Feuer EJ, Brown ML. Projections of the cost of cancer care in the United States: 2010-2020. Journal of the National Cancer Institute. Jan 19 2011;103(2):117-128.
- 3. Howlader N NA, Krapcho M, Garshell J, et al. SEER Cancer Statistics Review, 1975-2012. <u>http://seer.cancer.gov/csr/1975-2012/</u>
- Avalere Health, LLC. Total cost of cancer care by site of service: physician office vs outpatient hospital. Community Oncology Alliance website. <u>http://www.communityoncology.org/pdfs/avalere-cost-of-cancer-care-study.pdf</u>. Published March 2012.
- 5. Berkley Research Group. 340B growth and the impact on the oncology marketplace. Washington DC: Berkley Research group; 2015.
- Fitch K, Pyenson B. Site of service cost differences for Medicare patients receiving chemotherapy. Milliman Client Report. Oct 19,2011. Access at: <u>http://us.milliman.com/insight/health/Site-of-Service-Cost-Differences-for-Medicare-Patients-Receiving-Chemotherapy/</u>
- Fitch K, Iwasaki K, Pyenson B. Comparing episode of cancer care costs in different settings: An actuarial analysis of patients receiving chemotherapy. Milliman Client Report. August 29, 2013. http://us.milliman.com/uploadedFiles/insight/2013/comparingepisode-cancer-care.pdf
- 8. Hayes J, Hoverman JR, Brow ME et.al., Cost differential by site of service for cancer patients receiving chemotherapy. Am J Manag Care. 2015;21:e189-e196.
- 9. Tangka FK, Trogdon JG, Richardson LC et. al. Cancer treatment cost in the United States. Cancer. 201;116:3477-84.
- 10. US Food and Drug Administration (FDA). Hematology/Oncology (Cancer) Approvals & Safety Notifications. Updated 1/28/2016. Available at http://www.fda.gov/drugs/informationondrugs/approveddrugs/ucm279174.htm
- 11. Conti RM, Landrum MB, Jacobson M. The impact of provider consolidation on outpatient prescription drug-based cancer care spending. Issue brief. Health Care Cost Institute. February 2016:1-11. Available at http://www.healthcostinstitute.org/files/HCCI-Issue-Brief-Impact-of-Provider-Consolidation.pdf
- 12. Medicare Payment Advisory Commission (MedPAC). Report to Congress: Overview of the 340B Drug Pricing program. May 2015. Available http://www.medpac.gov/documents/reports/may-2015-report-to-the-congress-overview-of-the-340b-drug-pricing-program.pdf?sfvrsn=0
- 13. Bergsagel PL. Where we were, where we are, where we are going: progress in multiple myeloma. Am Soc Clin Oncol Educ Book. 2014:199-203.
- 14. Cunningham D, Atkin W, Lenz HJ, et al. Colorectal cancer. Lancet. 2010 Mar 20: 375(9719): 1030-47.
- 15. Carlson B. KRAS Testing: Optimizing Cancer Therapy. Personalized Medicine. Winter 2009-2010: 7-9.